# SOUTH BAY UROLOGY MEDICAL GROUP, INC.

**A DIVISION OF USSC** 

Joy G. Paul, MD, FRCS (C), FACS Garrett S. Matsunaga, MD Timothy Lesser, MD Shelby Morrisroe, MD 20911 Earl Street, Suite 140, Torrance, CA 90503
Phone 310.542.0199
Fax 310.542.4652
www.SouthBayUrology.com

## **BIOPSY OF THE PROSTATE**

A prostate biopsy is a procedure that will help your doctor to determine if you have prostate cancer or - in some cases - to monitor for progression of an existing prostate cancer. It is performed using ultrasound guidance via a probe that is placed in the rectum. The ultrasound will allow your doctor to visualize the prostate for any abnormalities, determine the size of the prostate, and to target the biopsy needle. It is a routine procedure usually done in the office or at a surgery center.

#### Before the procedure

- Allow approximately 60-90 minutes for the appointment; you may drive yourself
- No dietary restrictions or changes are required
- Take one Cipro 500 mg (or other antibiotic if allergic) the morning of the procedure
- Inform your doctor if you have a prosthetic implant that requires antibiotics prior to invasive procedures
- Use a Fleet's Enema (available in any pharmacy) approximately two hours prior to the procedure
- Be certain that you stop taking any medications that increase the chance of bleeding (see attached list)
   one week prior to the procedure

#### The procedure

You will be escorted into the examination room and placed on your left side on the examination table. The ultrasound technician will then give you an additional injection of an antibiotic and then introduce the ultrasound probe into the rectum. S/he will then determine the size of the prostate and assess for any abnormalities.

The doctor will then enter the room and inject local anesthetic around the area of the prostate to decrease the discomfort from the biopsies. Twelve needle biopsies will be taken. In certain situations, more or fewer samples will be taken. During the procedure, you may notice increasing discomfort in the prostate, bladder, or urethral area and/or a sensation that you have to urinate. This is normal. The procedure itself will take no more than 10 minutes.

### After the procedure

Take the second dose of Cipro 500 mg (or other antibiotic if allergic) the evening of the biopsy. It is expected to have hematuria (blood in the urine) and hematochezia (blood in the stool) for a few days and even up to a few weeks after the procedure. Hematospermia (blood in the semen) is not unusual for up to a few months after the biopsy. Take it easy for the first couple of days, though you may return to work the day after the biopsy.

### Please report the following conditions:

- Heavy bleeding which is worsening (with blood clots) either in the urine or stools
- Fever > 101<sup>0</sup>F
- Inability to urinate

As always, please do not hesitate to contact the office if any issues or questions arise

## SOUTH BAY UROLOGY MEDICAL GROUP, INC.

#### **A DIVISION OF USSC**

Joy G. Paul, MD, FRCS (C), FACS Garrett S. Matsunaga, MD Timothy Lesser, MD Shelby Morrisroe, MD 20911 Earl Street, Suite 140, Torrance, CA 90503 Phone 310.542.0199 Fax 310.542.4652 www.SouthBayUrology.com

# **Medications with Bleeding Risk**

#### Α

A.A. Compound
A.C.A. #4 Capsule
Acephens Suppositories
Advil
Aleve
Alka-Seltzer plus
Alka-Seltzer Plus Sinus Allergy
Alka-Seltzer Plus Cold and
Night Time Cold Medicine
Allerest Headache Strength

Anluprin Ampheno 1 Anacin Anacin Caps/Tablets Anacin Maximum Analyal

Amhralgen Tablets Amhralgen Amhropan

Anodynos

Arthritis Pain Formula Arthritis Strength Bufferin

ASA Compound Capsules
Ansemco #2 Tablets

ASA Enseals APAC Tablets APAP Tablets Aspercin

Aspir 10 (Vale)

**Azdone Tablets** 

Asphac G Tablets Ascritptin A/D Tablets Ascriptin Xstrength Tablets Asprabuf Aspergum

В

B.A.
Bayer Aspirin
Bayer Children's Aspirin
Bayer Decongestant Tablets
Bayer Maximum
Bayer Timed Release Aspirin

BC All clear

BC Tablets & Powder Bextra Buf Tablets

Buf Tablet Bufex Buff A

Buffadyne Lemon Buffaprin

Bufferin

Bufferin (Arthritis Strength) Buffering Xstrength

Buffets Buffinol C Cana (In Lay Tablets)

Celebrex Chewable Tablets Clinoril

Congesprin Chewables

Cope Coricidin Coricidin D

Coricidin Demilets (Children's)

Coricidin Medlets Cosprin Coumadin

CPZ Tablets Cystex

D

Damason
Datril
Darril 500
Dewitt Pills
Disalcid
Dasin Capsules
Doans Pills
Dolcin
Dolor

Dristan Duoprin Capsules Duoprin Syrup Duradyne Dularin Syrup

Ε

**Dynosal Tablets** 

Easprin
Ectrin
Emagrin
Empirin Analgesic
Empirin Compound
Empirin with Codeine
Encapin

Equagesic Tablets Excedrin Excedrin PM

**Excedrin APAP Tablets** 

**F** inol

Febrinol
Feldene
Femcaps
Fiorinal Caplets & Tablets
Fizzin Powder
Four Way Cold Tablets

G

Gavsals Tablets Gemnisyn (Rorer) Goody's Extra Strength Tablets Goody's Headache Powder Н

Hyalex Hipirn (Blane)

-

Ibuprofen Indocin

L

Liquiprin Solution Lortab ASA Tablets

M

Magneprin Magnaprin Arthritis Strength Meadache Measurin (Time Release) Meclomen Midol (Ibuprofen)

Midol (Ibuprofer Mobigestic

Momentum Muscle Backache

Ν

Nalfon Naprosyn Neocylate Neolin Nilain Nilprin 7 ½ Norwich ASA

Pabralate

Ρ

Pabirin
PAC
Pepto Bismol Tablets
Percodan
Percogesic
Presalin

Panodynes Analgesic

Q

R

**Quiet World Tablets** 

Rufen

SineAid

S

SAC Salatin Saleto SalFayne Capsules Salocol Saw-Palmetto Sinearest
SineOff
SK-Apap tablets
Soma Compound
St. Joseph Aspirin (children's)
St Joseph Cold (Tablets for child)
Stanback Tablets & Powder
Stanco
Supac
Synalgos DC Capsules

Т

Talwin Compound
Tempura Syrup & Drops
Tenol
Therapy Bayer ASA Caplets
Triagesic
Tn Pain Capletes
Tniaminicin
Triisate

U

Uracel 5 Ursinus

٧

Valadol Tablets & Liquid Valesin Vanequish Caplets Verin Viro-Med Tablets Vitamin E Vioxx

W

Weseprin Buffered

Ζ

Zocor Zorpin

# ANTIDIARRHEAL MEDICATIONS

(Beecham) Corrective Measure (Pfeiffer) Kaodene Non-Narcotic (Rexall) Pabinol with Paragoric (Norwich) Pepto Bismol